

# Fruit Crops Blueberries

## Blueberry Spray Program

| Timing                        | Pest        | Pesticide                                                                      | Amt/<br>100 gal | Remarks                   |
|-------------------------------|-------------|--------------------------------------------------------------------------------|-----------------|---------------------------|
| <b>Pre-Bloom Sprays</b>       |             |                                                                                |                 |                           |
| 1. Green Tip                  | Mummy Berry | Abound<br>Bravo or Equus<br>Captan<br>Captevate<br>Pristine<br>Switch<br>Ziram | See Mfg. label  |                           |
| 2. 7-14 days later            | Mummy Berry | Abound<br>Bravo or Equus<br>Captan<br>Captevate<br>Pristine<br>Switch<br>Ziram | See Mfg. label  |                           |
| <b>Bloom Sprays</b>           |             |                                                                                |                 |                           |
| 3. 10 - 20% Bloom             | Mummy Berry | Abound<br>Bravo or Equus<br>Captan<br>Captevate<br>Pristine<br>Switch<br>Ziram | See Mfg. label  |                           |
| 4. Full Bloom                 | Mummy Berry | Abound<br>Bravo or Equus<br>Captan<br>Captevate<br>Pristine<br>Switch<br>Ziram | See Mfg. label  |                           |
| <b>Petal Fall Sprays</b>      |             |                                                                                |                 |                           |
| 5. Immediately<br>After Bloom | Fruit Rots  | Cabrio<br>Captan<br>Captevate<br>Pristine<br>Rovral<br>Switch<br>Ziram         | See Mfg. label  | Rovral for Botrytis only. |

## Fruit Crops Blueberries

|                                |                          |                                                                       |                |                                               |
|--------------------------------|--------------------------|-----------------------------------------------------------------------|----------------|-----------------------------------------------|
| 6. 10-14 days later            | Fruit Rots<br>Leaf Spots | Cabrio<br>Captan<br>Captevat<br>Pristine<br>Rovral<br>Switch<br>Ziram | See Mfg. label | Rovral for Botrytis only.                     |
| 7. 10-14 days later            | Fruit Rots<br>Leaf Spots | Cabrio<br>Captan<br>Captevat<br>Pristine<br>Rovral<br>Switch<br>Ziram | See Mfg. label | Rovral for Botrytis only.                     |
| Fall Sprays                    | Bacterial Canker         | Fixed Copper (Kocide 2000,<br>Kocide DF, Nu-Cop 3L)                   | See Mfg. label | Make 2 sprays in fall 4 weeks<br>apart.       |
| Spring (before root<br>growth) | Phytophthora<br>Root Rot | Aliette or Helena Prophyt or<br>Phostrol<br>Ridomil Gold EC           | See Mfg. label | See manufacturer's label for<br>instructions. |